



TITLE:

睡眠時無呼吸症候群の心臓交感神経機能及びQT間隔のバラツキとレプチン分泌動態

AUTHOR(S):

陳, 和夫

CITATION:

陳, 和夫. 睡眠時無呼吸症候群の心臓交感神経機能及びQT間隔のバラツキとレプチン分泌動態. 2003

ISSUE DATE:

2003-05

URL:

<http://hdl.handle.net/2433/85211>

RIGHT:

学術雑誌掲載論文の抜き刷り、出版社に著作権許諾が得られていないため未掲載。

はじめに

閉塞性睡眠時無呼吸低呼吸症候群(obstructive sleep apnea-hypopnea syndrome:OSAHS)は本邦においても成人男子の数%以上にみられる頻度の高い症候群である。OSAHSは高血圧発症の独立因子であることは米国の大規模試験で最近明らかになったが、心・脳血管障害発症により予後が悪化するとも言われている。また、OSAHS70%以上は肥満を示す。

睡眠時無呼吸症候群の心臓交感神経機能 及びQT間隔のバラツキとレプチン分泌動態

本研究の目的は、睡眠時無呼吸症候群患者で治療前後に検討し、心臓交感神経機能とQT分散(QT dispersion)をOSAHS患者で治療前後に検討し、心臓交感神経機能との関連を検討すること、交感神経機能、睡眠に大きく関与するレプチンの分泌動態をOSAHS患者で治療前後に検討すること。

(13670598)

平成13年度～平成14年度科学研究費補助金(基盤研究C2)研究成果報告書

研究組織

研究代表者 陳 和夫(京都大学医学研究科 助教授)

研究分担者 三輪 理見(京都大学医学研究科 教授)

交付決定額

平成13年度 2,700千円

平成14年度 700千円

合計 3,400千円

平成15年5月

研究代表者 陳 和夫(京都大学医学研究科 助教授)

京 都 大 学 図 書 館



9810059501

附 属 図 書 館

はじめに

閉塞性睡眠時無呼吸低呼吸症候群(obstructive sleep apnea-hypopnea syndrome:OSAHS)は本邦においても成人男子の数%以上にみられる頻度の高い症候群である。OSAHS は高血圧発症の独立因子であることは米国の大規模試験で最近明らかになったが、心・脳血管障害発症により予後が悪化するとも言われている。また、OSAHS70%以上は肥満を示すが、OSAHS は肥満関連脂質代謝異常にも関与するとも言われている。

本研究の目的は、心臓死発症の独立因子である心電図上の QT 時間の最大値と最小値の差 QT dispersion を OSAHS 患者で治療前後に検討し、心臓交感神経機能との関連を検討すること、交感神経機能、脂質代謝に大きく関与するレプチンの分泌動態を OSAHS 患者で治療前後に検討することである。

研究組織

研究代表者 陳 和夫(京都大学医学研究科 助教授)

研究分担者 三嶋 理晃(京都大学医学研究科 教授)

交付決定額

平成 13 年度 2,700 千円

平成 14 年度 700 千円

総計 3,400 千円

陳和夫、中野敬雄、紀岡大代、市原研、八嶋茂光、大井元晴 閉塞性睡眠時無呼吸低呼吸症候群 OSAHS の時間的レプチンとの関連。第 42 回日本呼吸器学会総会、2002.4.5.仙台

上田和幸、谷口正幸、川口正、杉田敏子、赤木久恵、高橋典子、岡村雅子、大井元晴、中野敬雄、紀岡大代、市原研、八嶋茂光、大井元晴 OSAHS 症例に対する日中経皮的 CPAP による Nasal CPAP 導入の試み。第 27 回日本睡眠学会学術大会、2002.7.4.名古屋

陳和夫、三嶋理晃、中村敦樹、江川英夫、木内哲也、三嶋理晃、Collin E. Sullivan, D.P. 三嶋、三嶋理晃、市原研の急性呼吸不全症例に対する非侵襲的補助換気。第 38 回日本呼吸器学会、2002.10.17.東京

陳和夫、日本呼吸器学会、睡眠時無呼吸症候群-6 章、肥満と睡眠時無呼吸症候群、2000 195-200

研究発表

〈学会誌等〉

1. Chin K. Obstructive sleep apnea-hypopnea syndrome and multiple risk factors for cerebrovascular and cardiovascular disease. *Recent Res Devel Resp Critical Care Med* 2001;1: 95-100.
2. Shimizu K, Chin K, Nakamura T, Masuzaki H, Ogawa Y, Hosokawa R, Niimi A, Hattori N, Nohara R, Sasayama S, Nakao K, Mishima M, Nakamura T, Ohi M. Plasma leptin levels and cardiac sympathetic function in patients with obstructive sleep apnoea-hypopnoea syndrome. *Thorax* 2002;57: 429-434.
3. Chin K, Nakamura T, Takahashi K, Sumi K, Ogawa Y, Masuzaki H, Muro S, Hattori N, Matsumoto H, Niimi A, Chiba T, Nakao K, Mishima M, Ohi M, Nakamura T. Effects of obstructive sleep apnea syndrome on serum aminotransferase levels in obese patients. *American Journal of Medicine* 2003;114:370-376.
4. Hida W, Okabe S, Tatsumi K, Kimura H, Akashiba T, Chin K, Ohi M, Nakayama H, Satoh M, Kuriyama T. Nasal continuous positive airway pressure improves quality of life in obesity hypoventilation syndrome. *Sleep Breath* 2003;7:3-12.
5. Niimi A, Matsumoto H, Ueda T, Takemura M, Suzuki K, Tanaka E, Chin K, Mishima M, Amitani R. Impaired cough reflex in patients with recurrent pneumonia. *Thorax* 2003; 58:152-153.

〈口頭発表〉

- 1) Chin K, Shimizu K, Nakamura T, Mishima M, Ohi M, Nakamura T. Obstructive sleep apnea syndrome may be an aggravating factor in patients with non-alcoholic steatohepatitis with serum aspartate aminotransferase levels at or above 40 IU/L. 2001 International Conference ATS 2002, 5.23, 2001, San Francisco CA.
- 2) Chin K, Nakamura T, Mishima M, Nakamura T, Ohi M. Obstructive sleep apnea-hypopnea syndrome and multiple risk factors for cerebro and cardiovascular diseases. In Symposium: Cardiovascular morbidity in sleep apnoea syndrome: cellular and biochemical mechanisms. 16th Congress of the European Sleep Research Society, 3-7 June 2002, Reykjavik.
- 3) Chin K, Nakamura T, Miyaoka F, Muro S, Mishima M, Ohi M, Nakamura T. Hyperleptinemia and effects of obstructive sleep apnea-hypopnea syndrome on liver function in obesity. 98th International Conference ATS 2002, 5.19, 2002.
- 4) Nakamura T, Chin K, Shimizu K, Ohi M, Mishima M. Effects of nCPAP therapy on QT dispersion in patients with obstructive sleep apnea-hypopnea syndrome. The 7th Asian Pacific Society of Respirology, 2002, 10.26, Taipei.
- 5) 陳和夫, 中村敬哉, 宮岡史代, 室繁郎, 三嶋理晃, 大井元晴. 閉塞性睡眠時無呼吸低呼吸症候群(OSAHS)の肝障害とレプチンとの関連. 第 42 回日本呼吸器学会総会, 2002.4.5.仙台
- 6) 上田和幸, 谷口充孝, 江川功, 杉田淑子, 村木久恵, 茶園真紀子, 原田順子, 太田陽子, 中井直治, 角谷寛, 陳和夫, 大井元晴. 重症 OSAHS 症例に対する日中睡眠ポリグラフィによる Nasal CPAP 導入の試み. 第 27 回日本睡眠学会定期学術集会, 2002.7.4.仙台
- 7) 陳和夫, 上本伸二, 中村敬哉, 江川裕人, 木内哲也, 三嶋理晃, Collin E. Sullivan, 田中紘一, 中村孝志. 肝移植後の急性呼吸不全症例に対する非侵襲的補助換気. 第 38 回日本移植学会総会, 2002.10.19.東京

〈出版物〉

陳 和夫, 他. 日本評論社. 睡眠時呼吸障害-6 章, 肥満と睡眠時呼吸障害. 2002 195 pp93-100

ORIGINAL ARTICLE

Plasma leptin levels and cardiac sympathetic function in patients with obstructive sleep apnea-hypopnea syndrome

Nakamura T, Chin K, Shimizu K, Ohi M, Mishima M. Effects of nCPAP therapy on QT dispersion in patients with obstructive sleep apnea-hypopnea syndrome.

Abstract

Objective: To test the hypothesis that QT dispersion, which is defined as the longest QT interval minus the shortest QT interval, in patients with obstructive sleep apnea-hypopnea syndrome (OSAHS) changes with nasal continuous positive airway pressure (nCPAP) therapy.

Methods: We studied 48 patients with OSAHS [45 men and 3 women; age 45.9 ± 10.8 yr, mean \pm SD; body mass index (BMI) 30.2 ± 4.3 kg/m² and apnea-hypopnea index (AHI) 51.9 ± 18.5 /hr] who underwent polysomnography before nCPAP was started and on the first night of nCPAP. We excluded patients with cardiac disease or arrhythmia sufficiently severe to require treatment. Electrocardiograms (ECGs) were recorded in these patients for 30 seconds before, during, and after sleep, during each polysomnography, and in the morning about one month later to calculate ECG values including the QT interval. ECGs were also recorded in 26 age-matched normal subjects (24 men and 2 women; age 45.9 ± 13.4 yr and BMI 23.7 ± 2.2 kg/m²) in the morning to obtain normal ECG values. All data were subjected to nonparametric tests.

Results: Morning values for the heart rate-corrected QT interval (QTcI) and QTc dispersion (QTcD) did not differ in OSAHS patients from normal subjects before, after one night or after one month of nCPAP therapy; Morning values for QTcI and QTcD were 423.6 ± 26.2 mm^{1/2} and 60.1 ± 12.8 mm^{1/2} before nCPAP, 421.4 ± 24.6 and 57.0 ± 12.3 after one night nCPAP, and 426.6 ± 28.1 and 57.6 ± 12.5 after one month nCPAP, respectively, in OSAHS patients vs 427.6 ± 22.6 and 59.5 ± 16.0 , respectively, in normal subjects. QTcI during sleep did not change with one night nCPAP (434.4 ± 38.3 mm^{1/2} before nCPAP vs 426.3 ± 24.3 with one night nCPAP). Meanwhile, QTcD during sleep significantly decreased with one night nCPAP (65.0 ± 14.5 mm^{1/2} before nCPAP vs 50.5 ± 11.3 with one night nCPAP, $p < 0.0001$). QTcD during sleep before nCPAP was significantly correlated with mean arterial oxygen saturation (SaO₂) ($r = -0.425$, $p = 0.004$) and the percentage of time that SaO₂ was below 90% ($r = 0.347$, $p = 0.017$) during sleep before nCPAP. The reduction in QTcD during sleep with one night nCPAP was significantly correlated with the reduction in AHI ($r = 0.366$, $p = 0.012$).

Conclusions: The present study suggests that nCPAP therapy decreases nocturnal myocardial electrical instability in patients with OSAHS even in the absence of cardiac disease or severe arrhythmia.

Obstructive sleep apnea-hypopnea syndrome (OSAHS) is a common disorder characterized by recurrent episodes of partial or complete upper airway obstruction during sleep, leading to intermittent hypoxemia and sleep fragmentation. OSAHS is associated with various cardiovascular complications, including hypertension, coronary artery disease, and stroke. The pathogenesis of OSAHS involves anatomical and physiological factors, and treatment options include continuous positive airway pressure (CPAP) therapy, oral appliances, and surgery. This study focuses on the effects of nasal continuous positive airway pressure (nCPAP) therapy on QT dispersion in patients with OSAHS. QT dispersion is a measure of the variability of the QT interval across different leads, which is thought to reflect myocardial electrical instability. The study aims to test the hypothesis that nCPAP therapy reduces QT dispersion in OSAHS patients, even in the absence of cardiac disease or severe arrhythmia. The study included 48 OSAHS patients and 26 age-matched normal subjects. ECGs were recorded at multiple time points, and nonparametric tests were used for data analysis. The results show that nCPAP therapy significantly reduced QTcD during sleep in OSAHS patients, and this reduction was correlated with improvements in oxygen saturation and AHI. These findings suggest that nCPAP therapy may have beneficial effects on myocardial electrical stability in OSAHS patients.